

ON FANO VARIETIES WITH LARGE PSEUDO-INDEX

JIUN-CHENG CHEN

ABSTRACT. Let X be a Fano variety with at worst isolated quotient singularities. Our result asserts that if $C \cdot (-K_X) > \max\{\frac{n}{2} + 1, \frac{2n}{3}\}$ for every curve $C \subset X$, then $\rho_X = 1$.

1. INTRODUCTION

We work over the field of complex numbers $\mathbb{C}$. Let X be a smooth Fano variety of dimension n . The index r_X is defined as

$$r_X = \max\{m \in \mathbb{N} \mid K_X = mL \text{ for some line bundle } L\}.$$

The pseudo-index i_X is defined as

$$i_X = \min\{m \in \mathbb{N} \mid C \cdot (-K_X) = m \text{ for some rational curve } C \subset X\}.$$

Clearly, we have $i_X \geq r_X$. Denote by ρ_X the Picard number of X . In 1988, Mukai [Mu88] proposed the following conjecture:

Conjecture 1.1 (Mukai Conjecture). *Let X be a smooth Fano variety of dimension n . Then $\rho_X \cdot (r_X - 1) \leq n$.*

A generalized version of this conjecture can be stated as follows:

Conjecture 1.2 (Generalized Mukai Conjecture). *Let X be a smooth Fano variety of dimension n . Then (1) $\rho_X \cdot (i_X - 1) \leq n$, and (2) equality holds if and only if $X \cong (\mathbb{P}^{i_X})^{\rho_X}$.*

For each integer $k \geq 1$, we can consider the following conjecture.

Conjecture 1.3 (Generalized Mukai Conjecture _{k}). *Let X be a smooth Fano variety of dimension n .*

- (1) *Suppose that $i_X > \frac{n}{k} + 1$. Then $\rho_X \leq k - 1$.*
- (2) *Suppose that $i_X \geq \frac{n}{k} + 1$ and $\rho_X = k$. Then $X \cong (\mathbb{P}^{\frac{n}{k}})^k$.*

Clearly, Conjecture 1.3 for all k implies Conjecture 1.2.

When $k = 1$, the first part of Conjecture 1.3 implies that if $i_X > n + 1$ then $\rho_X \leq 0$. Since the Picard number of any projective variety is always at least 1, it implies that $i_X \leq n + 1$ for any smooth Fano variety of dimension

Date: October 28, 2009.

Research of J.-C. C supported in part by National Center of Theoretical Science in Hsinchu, Taiwan, National Science Council, Taiwan, and by the Golden Jade Fellowship of Kenda Foundation.

n . This consequence follows from Mori's famous bend and break result. The second part of Conjecture 1.3 follows from the characterization result of projective spaces by Cho, Miyaoka and Shepherd-Barron [CMSB02] and Kebekus [Ke01]. For $k = 2$, Wiśniewski [Wi90] proved that if $i_X > \frac{n}{2} + 1$ then $\rho_X = 1$.

For $k = 3$, there are related results due to Bonavero, Casagrande, Debarre and Druel [BCDD03], and Andreatta, Chierici, and Occhetta [ACO04]. We state their precise results here. Bonavero, Casagrande, Debarre and Druel [BCDD03] proved Conjecture 1.2 in the following situations: (1) X has dimension 4, (2) X is a toric variety of pseudo-index $i_X \geq \frac{n}{3} + 1$ or of dimension at most 7. Andreatta, Chierici, and Occhetta [ACO04] proved the generalized Mukai conjecture assuming either (1) $i_X \geq \frac{n}{3} + 1$ and X has a fiber type extremal contraction, (2) $i_X \geq \frac{n}{3} + 1$ and X has not small extremal contractions, or (3) $\dim X = 5$.

Now consider the case when X is not smooth. Even the $k = 1$ case, i.e. $i_X \leq n + 1$, is not known. When X has only quotient singularities or isolated LCIQ singularities, Tseng and I proved that $i_X \leq n + 1$ [CT09]. In the same paper, we also proved if X is log terminal and $\dim X = 3$ then $i_X \leq 3 + 1$.

In [CT07], we proved that (1) if X has at worst isolated LCIQ singularities and $i_X \geq n + 1$ then $X \cong \mathbb{P}^n$, and (2) if X a normal $\mathbb{Q}$ -factorial Fano threefold and $i_X \geq 4$ then $X \cong \mathbb{P}^3$. In [Ch06], we proved that if X has at worst quotient singularities and $i_X \geq n + 1$ then $X \cong \mathbb{P}^n$.

In this note, we would like to study the $k = 2$ case when the variety X has only isolated quotient singularities. The following theorem is the main result of this note.

Theorem 1.4. *Let X be a Fano variety of dimension n with at worst isolated quotient singularities. If $i_X > \max\{\frac{n}{2} + 1, \frac{2n}{3}\}$, then $\rho_X = 1$.*

Consider a smooth Fano variety X . There are a lot of rational curves on X and this fact is important in studying the geometry of X . The deformation theory of rational curves is well-known and has many important consequences. When X is not smooth, the situation is quite different. The dimension of deformations (of rational curves) tends to be smaller. It is also possible that all rational curves pass through a singular point. These obstacles make the study of deformations of rational curves on singular Fano varieties quite difficult. When X has only quotient singularities, one natural approach is to consider the covering stack $\mathcal{X}$ and study "rational curves" on the covering stack $\mathcal{X}$. Pushing forward any family of curves on $\mathcal{X}$ to X yields a family of curves on X . If we can obtain a nice family of rational curves on $\mathcal{X}$, we have a nice family of rational curves on X for free. Let $g : C \rightarrow \mathcal{X}$ be a rational curve. The lower bound on the dimension of the deformation space of $g : C \rightarrow \mathcal{X}$ can be computed as in the smooth case. Roughly speaking, the deformation theory does not see the twisted points on the target $\mathcal{X}$. So far, everything seems work. But there are several difficulties for this approach. One problem is that when we have a curve $g : C \rightarrow X$ it may not be possible

to lift it to $\tilde{g} : C \rightarrow \mathcal{X}$. To get a lifting, one has to add twisted points on the source curve C [AV02]. Therefore, we have to study the deformations of twisted curves into $\mathcal{X}$. Unfortunately, the presence of twisted points on the source curve will lower the dimension of the deformation space (due to the age terms, see Lemma 2.2 for the precise formula). The second bad news is that even if we start with a curve (without twisted points) $g : C \rightarrow \mathcal{X}$, we may have to consider twisted points after applying bend and break to obtain curves of smaller degree.

In [CT09] and [CT07], we developed several techniques to handle the problems caused by the presence of twisted points. One of our main results in [CT09] is that if we start with a curve (in an extremal ray) without twisted point then we can find a twisted curve with at most one twisted point in the same extremal ray and bound the degree at the same time. Twisted curves with only one twisted point are good enough in many cases when we try to apply bend and break type arguments.

The starting point of this work is a very naive dimension count. Suppose that $\rho_X \geq 2$. We can find two extremal classes α and β . Lemma 2.8 guarantees that we can find twisted curves $f_1 : \mathcal{C}_1 \rightarrow \mathcal{X}$ and $f_2 : \mathcal{C}_2 \rightarrow \mathcal{X}$ whose classes are α and β respectively. For $i = 1, 2$, the twisted curve $\mathcal{C}_i$ has at most one twisted point and $\mathcal{C}_i \cdot -K_X \leq n + 1$. For simplicity, assume both $\mathcal{C}_1$ and $\mathcal{C}_2$ do have one twisted point. Let $V_1 \subset \mathcal{X}$ be the subset swept by the deformation of $f_1(\mathcal{C}_1) \subset \mathcal{X}$ and $V_2 \subset \mathcal{X}$ the subset swept by the deformation of $f_2(\mathcal{C}_2) \subset \mathcal{X}$. Since the twisted points on $\mathcal{X}$ are isolated, the image of the twisted point on $\mathcal{C}_1$ does not deform. Therefore every deformation of $f_1(\mathcal{C}_1)$ has to pass through the specified point. Similarly, every deformation of $f_2(\mathcal{C}_2)$ also has to pass through another specified point. Lemma 2.9 implies that these two sets should not intersect. It is easy to see $\dim V_1 \geq i_X - 1$ and $\dim V_2 \geq i_X - 1$. We "expect" them to intersect if $2i_X - 2 - n > 0$, i.e. $i_X > \frac{n}{2} + 1$. Therefore, the assumption $\rho_X \geq 2$ is wrong whenever $i_X > \frac{n}{2} + 1$. Of course, this "argument" is very problematic since we can have two disjoint divisors in a variety.

To remedy this incomplete argument, we consider a covering family $\mathcal{C} \rightarrow \mathcal{X}$ of twisted stable maps. Note that there may be many twisted points on a general source curve $\mathcal{C}_t$.

We only sketch the argument for the worst case, i.e. the source curve $\mathcal{C}_t$ has at least two twisted points. In general, the presence of twisted points on the source curve causes troubles. However, when X_{sing} is isolated, we can play the presence of many twisted points to our advantage via bend and break type arguments: the images of these twisted points can not move; Mori's bend and break result then implies this covering family will degenerate somewhere.

Assume that $\mathcal{C}_t$ has at least 2 twisted points. Using a bend and break type argument, we can show that there is a family of unsplit curves such that the locus of this family contains at least a divisor (see Proposition 3.3 for more details). Denote this divisor by D . Once we have such a family

of curves, we find an extremal contraction, denoted by $\phi : X \rightarrow Y$, whose exceptional set contains the divisor D . We will show that Y is a point. This contraction is either a divisorial contraction or a contraction of fiber type. The contraction $\phi : X \rightarrow Y$ has the following property: $C \cdot D > 0$ where C is any curve contracted by $\phi : X \rightarrow Y$.

If the contraction $\phi : X \rightarrow Y$ is divisorial, then the inequality $C \cdot D > 0$ gives a contradiction by a standard argument.

If the contraction is of fiber type and $\dim Y > 0$, then we can find a closed subset $Z \subset X$ such that $\dim Z > \dim Y$ and $Z \rightarrow Y$ is finite. This is again a contradiction (see Lemma 3.1 for more details).

The rest of this paper is organized as follows: In Section 2, we recall basic definitions on twisted curves and twisted stable maps. We also recall Lemma 7.1 from [ACO04] in Section 2. The main theorem is proved in Section 3.

ACKNOWLEDGEMENTS

The author likes to thank Dan Abramovich, Lawrence Ein and Hsian-Hua Tseng for helpful discussions and valuable suggestions. The author also likes to thank the referee for his valuable comments and suggestions.

2. TWISTED CURVES

We recall some basic facts on twisted curves in this section. All results in this section are not new; we recall these facts for the reader's convenience.

Twisted curves appear naturally in compactifying stable maps into a proper Deligne-Mumford stack [AV02]. Roughly speaking, twisted curves are nodal curves having certain stack structures étale locally near nodes (and, for pointed curves, marked points). For the precise definition, see [AV02], Definition 4.1.2.

Let $\mathcal{C}$ be a twisted curve and C its coarse moduli space.

2.0.1. Nodes. For a positive integer r , let μ_r denote the cyclic group of r -th roots of unity. Étale locally near a node, a twisted curve $\mathcal{C}$ is isomorphic to the stack quotient $[U/\mu_r]$ of the nodal curve $U = \{xy = f(t)\}$ by the following action of μ_r :

$$(x, y) \mapsto (\zeta_r x, \zeta_r^{-1} y),$$

where ζ_r is a primitive r -th root of unity. Étale locally near this node, the coarse curve C is isomorphic to the schematic quotient U/μ_r .

2.0.2. Markings. Étale locally near a marked point, $\mathcal{C}$ is isomorphic to the stack quotient $[U/\mu_r]$. Here U is a smooth curve with local coordinate z defining the marked point, and the μ_r -action is defined by

$$z \mapsto \zeta_r z.$$

Near this marked point the coarse curve is the schematic quotient U/μ_r .

2.1. Twisted stable maps.

Definition 2.1. A twisted n -pointed stable map of genus g and degree d over a scheme S consists of the following data (see [AV02], Definition 4.3.1):

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{f} & \mathcal{X} \\ \pi_{\mathcal{C}} \downarrow & & \downarrow \pi \\ C & \xrightarrow{\bar{f}} & X \\ \downarrow & & \\ S & & \end{array}$$

along with n closed substacks $\Sigma_i \subset \mathcal{C}$ such that

- (1) $\mathcal{C}$ is a twisted nodal n -pointed curve over S (see [AV02], Definition 4.1.2),
- (2) $f : \mathcal{C} \rightarrow \mathcal{X}$ is representable,
- (3) Σ_i is an étale gerbe over S , for $i = 1, \dots, n$, and
- (4) the map $\bar{f} : (C, \{p_i\}) \rightarrow X$ between coarse moduli spaces induced from f is a stable n -pointed map of degree d in the usual sense.

A twisted map $f : \mathcal{C} \rightarrow \mathcal{X}$ is stable if and only if for every irreducible component $\mathcal{C}_i \subset \mathcal{C}$, one of the following cases holds:

- (1) $f|_{\mathcal{C}_i}$ is nonconstant,
- (2) $f|_{\mathcal{C}_i}$ is constant, and $\mathcal{C}_i$ is of genus at least 2,
- (3) $f|_{\mathcal{C}_i}$ is constant, $\mathcal{C}_i$ is of genus 1, and there is at least one special points on $\mathcal{C}_i$,
- (4) $f|_{\mathcal{C}_i}$ is constant, $\mathcal{C}_i$ is of genus 0, and there are at least three special points on $\mathcal{C}_i$.

In particular, a nonconstant representable morphism from a smooth twisted curve to $\mathcal{X}$ is stable.

We say a twisted stable map $\mathcal{C} \rightarrow \mathcal{X}$ is rational if the coarse moduli space C of $\mathcal{C}$ is rational.

We will use the following two basic facts on twisted stable maps frequently in this note:

(1) Let $f : \mathcal{C} \rightarrow \mathcal{X}$ be a twisted stable map. Then the image of any twisted point on $\mathcal{C}$ has to be a twisted point on $\mathcal{X}$.

(2) Consider a family of rational twisted stable map $F : \mathcal{C} \rightarrow \mathcal{X}$ over an irreducible projective scheme T . If this family does not degenerate, i.e. (i) $\mathcal{C}_t$ is irreducible for every $t \in T$ and (ii) $F_t : \mathcal{C}_t \rightarrow \mathcal{X}$ is birational to its image for every t , then the number of twisted points on $\mathcal{C}_t$ is constant.

Fix an ample $\mathbb{Q}$ -line bundle H on X . Let $\mathcal{K}_{g,n}(\mathcal{X}, d)$ be the category of twisted n -pointed stable maps to $\mathcal{X}$ of genus g and degree d with respect to

the pull-back of H . We will set $H = -K_X$ when X is a Fano variety with at worst quotient singularities. It is known that $\mathcal{K}_{g,n}(\mathcal{X}, d)$ is a proper Deligne-Mumford stack with projective coarse moduli space denoted by $K_{g,n}(\mathcal{X}, d)$ [AV02]. Let $\beta \in H_2(X)$ be a homology class. The space of twisted n -pointed stable maps $f : \mathcal{C} \rightarrow \mathcal{X}$ of genus g and homology class $(\pi \circ f)_*[\mathcal{C}] = \beta$ is denoted by $\mathcal{K}_{g,n}(\mathcal{X}, \beta)$. This stack is also proper [AV02].

2.2. Morphism space from a twisted curve to a Deligne-Mumford stack. In this paper, we use both the stack of twisted stable maps and the morphism space from $\mathcal{C}$ to $\mathcal{X}$. Roughly speaking, an element in the morphism space $Mor(\mathcal{C}, \mathcal{X})$ is a twisted stable map together with a parametrization on the source curve $\mathcal{C}$. Let $\Sigma \subset \mathcal{C}$ be the set of twisted points and $B \subset \mathcal{C}$ a finite set of points (twisted or untwisted). Let $f : \mathcal{C} \rightarrow \mathcal{X}$ be a representable morphism. When $\mathcal{X}$ is smooth, we have a lower bound on the dimension of $Mor(\mathcal{C}, \mathcal{X}; f|_B)$ near the morphism $[f]$.

Lemma 2.2 (= [CT09] Lemma 4.4).

$$\dim_{[f]} Mor(\mathcal{C}, \mathcal{X}; f|_B) \geq -\mathcal{C} \cdot K_{\mathcal{X}} + n[\chi(\mathcal{O}_{\mathcal{C}}) - Card(B)] - \sum_{x \in \Sigma \setminus B} age(f^* T_{\mathcal{X}}, x).$$

Remark 2.3. For each twisted point $x \in \mathcal{C}$, the contribution from the age term is strictly less than $\dim X = n$ [CT09].

Consider

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{X} \\ g \downarrow & & \\ T & & \end{array}$$

a family of twisted stable maps. We are not only interested in the dimension of T but also the dimension of $F(\mathcal{C})$. In general, information on the dimension of T (the parametric space) does not yield much information on the dimension of $F(\mathcal{C})$ (the locus swept by the curves $\{\mathcal{C}_t | t \in T\}$). The main problem is that for any given point x there may be a positive dimensional family of curves passing through x . For a special class of curves, this problem disappears.

Definition 2.4. Let $C \subset X$ be a curve. We say C is *unsplit* if the homology class $[C] \in H_2(X)$ can not be written as a non-trivial positive integral sum of curve classes, i.e. if $[C] = \sum_{i=1}^k a_i [C_i]$ ($a_i \in \mathbb{N}_{>0}$) then $k = 1$ and $[C_1] = [C]$.

Definition 2.5. *Let*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{X} \\ \pi_{\mathcal{C}} \downarrow & & \downarrow \pi \\ C & \xrightarrow{\bar{F}} & X \\ \bar{G} \downarrow & & \\ T & & \end{array}$$

be a family of twisted stable maps. We say it is an unsplit family if F_t is birational to its image and $(\pi \circ F_t)(\mathcal{C}_t) \subset X$ is unsplit for every $t \in T$.

When the family is unsplit, we have the following bound:

Lemma 2.6. *Let*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{X} \\ \pi_{\mathcal{C}} \downarrow & & \downarrow \pi \\ C & \xrightarrow{\bar{F}} & X \\ \bar{G} \downarrow & & \\ T & & \end{array}$$

be an unsplit family of twisted 1-pointed stable maps. Assume the image of the marked point is fixed. Let $d = \mathcal{C}_t \cdot -K_{\mathcal{X}}$ be the anti-canonical degree. Then

$$\dim F(\mathcal{C}) \geq d - 1$$

Proof. This lemma follows easily from Mori's bend and break, Lemma 2.2, Remark 2.3 and the fact that $\dim \text{Aut}(\mathbb{P}^1, 0) = 2$. $\square$

2.3. Covering Stack and Lifting. We start with the notation of the covering stack $\mathcal{X}$.

Notation 2.7. *Let X be a normal projective variety with at worst quotient singularities. Fix a natural number $r \in \mathbb{N}_{>0}$ such that rK_X is Cartier. Fix a proper smooth Deligne-Mumford stack $\pi : \mathcal{X} \rightarrow X$ such that X is a coarse moduli space of $\mathcal{X}$ and π is an isomorphism over $X_{\text{reg}} = X \setminus X_{\text{sing}}$. Note that $K_{\mathcal{X}} = \pi^* K_X$ and*

$$\mathcal{C} \cdot K_{\mathcal{X}} = C \cdot K_X$$

for any (twisted) curve $\mathcal{C} \rightarrow \mathcal{X}$ with coarse curve C [CT09].

Let C be a smooth irreducible curve and $\bar{f} : C \rightarrow X$ a morphism. In general, it is not possible to “lift” the map $\bar{f} : C \rightarrow X$ to a map $C \rightarrow \mathcal{X}$. However, we can endow an orbicurve structure on C and lift $C \rightarrow X$ to $\mathcal{C} \rightarrow \mathcal{X}$ (by Lemma 7.2.5 [AV02]).

An arbitrary lifting is not very useful since the dimension lower bound obtained from Lemma 2.2 will be too small. One would like to have a lifting

with as few twisted points as possible. We do not know how to achieve this goal in general. If the class of $C \subset X$ is in an extremal ray, we are able to find a nice lifting as in the next lemma.

Lemma 2.8 (Proposition 3.2 [CT07]). *Notation as in Notation 2.7. Let $R \subset \overline{NE}(X)$ be any K_X -negative extremal ray. Then there exists a twisted rational curve $f : \mathcal{C} \rightarrow \mathcal{X}$ such that (1) $\mathcal{C}$ has at most one twisted point, (2) the intersection number $\mathcal{C} \cdot (-K_{\mathcal{X}}) \leq n + 1$, and (3) $(\pi \circ f)_*[C] \in R$.*

2.4. Twisted curves with at most 1 twisted point. Assume that X is a normal projective variety of dimension n with at worst isolated quotient singularities. Let $f : \mathcal{C} \rightarrow \mathcal{X}$ be an unsplit twisted stable map of genus 0 and homology class $\alpha \in H_2(X)$. Assume that the source curve $\mathcal{C}$ has at most 1 twisted point.

If $\mathcal{C}$ has one twisted point, denoted by $\infty \in \mathcal{C}$, take an irreducible component $\mathcal{M} \subset \mathcal{K}_{0,1}(\mathcal{X}, \alpha)$ which contains $[f]$. Take the universal family of twisted stable maps over $\mathcal{M}$. Note that the image of ∞ is fixed since X has only isolated singularities.

If $\mathcal{C}$ does not have any twisted point, i.e. $\mathcal{C} \cong \mathbb{P}^1$, pick any point, denoted by ∞ again. Consider the curve $f : \mathcal{C} \rightarrow \mathcal{X}$ with the marked point $\infty \in \mathcal{C}$ as an element of $\mathcal{K}_{0,1}(\mathcal{X}, \alpha)$. We abuse the notation and still denote this element in $\mathcal{K}_{0,1}(\mathcal{X}, \alpha)$ by $[f]$. Consider an irreducible component $\mathcal{M} \subset \mathcal{K}_{0,1}(\mathcal{X}, \alpha, f|_{\infty})$ which contains $[f]$. Take the universal family of twisted stable maps over $\mathcal{M}$.

In either case, we obtain a family of twisted stable maps such that the image of the marked point is fixed.

2.5. Bounding Picard Number. Let X be a normal projective variety and

$$\begin{array}{ccc} C & \xrightarrow{F} & X \\ G \downarrow & & \\ V & & \end{array}$$

be a family of stable maps of genus 0 over an irreducible scheme V .

We denote by $Locus(V)$ the image of C in X . Let $p \in X$ be a (closed) point. Define $Locus(V)_p := F(G^{-1}(G(F^{-1}(p))))$, i.e. the set of points $x \in X$ such that there is a $v_0 \in V$ satisfying $x \in F(C_{v_0})$ and $p \in F(C_{v_0})$.

The following lemma is a special case of Lemma 5.1 in [ACO04].

Lemma 2.9. *Let X be a normal projective variety, $p \in X$ any point, and V an unsplit family of rational curves. Also assume that V is irreducible and projective. Then $Locus(V)_p$ is closed and every curve contained in $Locus(V)_p$ is numerically equivalent to μC_V , where C_V is the class of $F(C_v)$ for $v \in V$ general and $\mu \geq 0$.*

2.6. Covering family of rational curves. The following lemma is an easy consequence of Kawamata's famous result [Ka91]:

Lemma 2.10. *Let X be a Fano variety of dimension n with at worst quotient singularities. Then there is a covering family of rational curves with $-K_X$ -degree at most $2n$.*

We have the following stack version of Lemma 2.10.

Lemma 2.11. *Let X be a Fano variety of dimension n . Assume that X has at worst quotient singularities. Let $\mathcal{X} \rightarrow X$ be the smooth covering stack as in Notation 2.7. Then there is a covering family of twisted rational curves with $-K_{\mathcal{X}}$ -degree at most $2n$.*

Proof. Let r be the natural number such that rK_X is Cartier as in Notation 2.7 and $d > 0$ be any rational number such that $rd \in \mathbb{N}_{>0}$. Consider the morphisms $\coprod_{k \in \mathbb{N}} \mathcal{K}_{0,k}(\mathcal{X}, d) \rightarrow \coprod_{k \in \mathbb{N}} \mathcal{K}_{0,k}(X, d)$ (forgetting the stack structures) and $\coprod_{k \in \mathbb{N}} \mathcal{K}_{0,k}(X, d) \rightarrow \mathcal{K}_{0,0}(X, d)$ (forgetting marked points and then stabilizing). Note that the composition $\coprod_{k \in \mathbb{N}} \mathcal{K}_{0,k}(\mathcal{X}, d) \rightarrow \mathcal{K}_{0,0}(X, d)$ is surjective since we can always lift a stable map $C \rightarrow X$ to $\mathcal{X}$ by adding suitable stack structures on finitely many points on C . Consider the universal family $F_{k,d} : \mathcal{U}_{k,d} \rightarrow \mathcal{X}$ over $\mathcal{K}_{0,k}(\mathcal{X}, d)$. It follows that

$$\cup_{\{(k,d) | k \in \mathbb{N}, d \leq 2n, rd \in \mathbb{N}\}} (\pi \circ F_{k,d})(\mathcal{U}_{k,d}) = X$$

since X is covered by rational curves with degree at most $2n$. Since X is Noetherian and irreducible, we have $(\pi \circ F_{k,d})(\mathcal{U}_{k,d}) = X$ for some (k, d) .

Since $\mathcal{K}_{0,k}(\mathcal{X}, d)$ is a proper Deligne-Mumford stack with projective coarse moduli space, we can find an étale surjective morphism $V \rightarrow \mathcal{K}_{0,k}(\mathcal{X}, d)$ such that V is projective.

Pulling back the universal family of twisted stable maps over $\mathcal{K}_{0,k}(\mathcal{X}, d)$ to V yields a covering family of twisted stable maps over V .

Taking a suitable irreducible component of V , we obtain a covering family over an irreducible projective scheme.

□

3. PROOF OF THEOREM 1.4

We will assume that X is a Fano variety of dimension n with at worst isolated quotient singularities throughout this section. Let $\pi : \mathcal{X} \rightarrow X$ be the covering stack of X as in Notation 2.7.

We start with the following important lemma.

Lemma 3.1. *Consider*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{X} \\ \pi_{\mathcal{C}} \downarrow & & \downarrow \pi \\ C & \xrightarrow{\bar{F}} & X \\ \bar{G} \downarrow & & \\ T & & \end{array}$$

an unsplit family of twisted stable maps over an irreducible projective scheme T . Assume that (1) the source curve $\mathcal{C}_t$ has at most 1 twisted point for a general $t \in T$, (2) the image $\bar{F}(C)$ contains a divisor in X and (3) $i_X > \frac{n}{2} + 1$. Then $\rho_X = 1$.

Proof. First note that $\mathcal{C}_t$ has at most 1 twisted point for any $t \in T$ since the family is unsplit. Consider the closed subset $V = \bar{F}(C)$, the image of the family of twisted stable maps in X . Since V is the image of an irreducible scheme under a projective morphism, it is closed and irreducible. Endow the reduced scheme structure on V . If $V = X$, take D be any irreducible divisor on X . If V is a divisor, take D be any prime divisor contained in V .

Pick a general $t \in T$ and consider the twisted stable map $F_t : \mathcal{C}_t \rightarrow \mathcal{X}$. Set $\beta = (\pi \circ F_t)_*[\mathcal{C}_t] = \bar{F}_t * [C_t] \in H_2(X)$.

Recall that X is $\mathbb{Q}$ -factorial. Let E be any curve which intersects D but is not contained in D . It is clear that $E \cdot D > 0$. Write $[E] = \sum a_i [E_i]$ as a positive combination of extremal rays. We have $E_i \cdot D > 0$ for some i . May assume that $E_1 \cdot D > 0$. Let $\alpha := [E_1]$. By Lemma 2.8, we can find a twisted rational curve $f : \mathcal{E} \rightarrow \mathcal{X}$ with at most one twisted point such that $(\pi \circ f)_*[\mathcal{E}] \in \mathbb{R}_{\geq 0} \alpha$ and $\mathcal{E} \cdot f^*(-K_{\mathcal{X}}) \leq n + 1$. Note that $f(\mathcal{E})$ is unsplit since $i_X > \frac{n}{2} + 1$.

Note that $\mathcal{E} \cdot (\pi \circ f)^*D > 0$ and therefore $f(\mathcal{E})$ meets $\pi^{-1}(D)$. Replacing α by the class $(\pi \circ f)(\mathcal{E})$ if necessary (they are two classes in the same ray), we may assume that $[(\pi \circ f)(\mathcal{E})] = \alpha \in H_2(X)$.

As in Section 2.4, we can view $f : \mathcal{E} \rightarrow \mathcal{X}$ as an element of $\mathcal{K}_{0,1}(\mathcal{X}, \alpha)$. Consider an irreducible component $[f] \in \mathcal{M} \subset \mathcal{K}_{0,1}(\mathcal{X}, \alpha)$ and its universal family of twisted 1-pointed stable maps as in Section 2.4. Denote by W_1 the image of this family in X . Note that for this family of twisted 1-pointed stable maps the image of the marked point is fixed.

By Lemma 2.9, it follows that for any curve $C \subset W_1$, its class $[C]$ lies in $\mathbb{R}_{\geq 0} \alpha$. We also have $\dim W_1 \geq \mathcal{E} \cdot (-K_{\mathcal{X}}) - 1 > \frac{n}{2}$ by Lemma 2.6.

We proceed by showing that the classes α and β lie in the same extremal ray. We then show that $\mathbb{R}_{\geq 0} \alpha$ is the only extremal ray.

Divide into two cases: (1) the source curve $\mathcal{C}_t$ for a general $t \in T$ does have a twisted point, and (2) the source curve $\mathcal{C}_t$ for a general $t \in T$ does not have any twisted point, i.e. $\mathcal{C} = C$ is a scheme.

Case (1): The source curve $\mathcal{C}_t$ has one twisted point.

Note that D is covered by a family of rational curves through a specified point (the image of the twisted point). By Lemma 2.9 again, it follows that for any curve $C \subset D$, its class $[C]$ lies in $\mathbb{R}_{\geq 0} \beta$. Consider $W_1 \cap D$. Since the intersection is non-empty, any irreducible component of $W_1 \cap D$ has dimension at least $\dim W_1 + \dim D - n \geq 1$. Therefore, we can find a curve $C_1 \subset D$ and $C_1 \subset W_1$. It follows that $\mathbb{R}_{\geq 0} \alpha = \mathbb{R}_{\geq 0} \beta$.

Case (2): The source curve $\mathcal{C}_t$ has no twisted point.

Pick any $t_0 \in T$ such that W_1 meets $(\pi \circ F_{t_0})(\mathcal{C}_{t_0})$. Pick any point on $\mathcal{C}_{t_0}$, say ∞ , and consider the morphism $F_{t_0} : \mathcal{C}_{t_0} \rightarrow \mathcal{X}$ with the marked point $\infty \in \mathcal{C}_{t_0}$ as a twisted 1-pointed stable map as in Section 2.4. We still denote this element by $[F_{t_0}]$. Let $\mathcal{M} \subset \mathcal{K}_{0,1}(\mathcal{X}, \beta, F_{t_0}|\infty)$ be an irreducible component which contains $[F_{t_0}]$. Consider the universal family of twisted 1-pointed stable maps over $\mathcal{M}$ and the image of this family in X . Denote the image by W_2 . Note that for any curve $C \subset W_2$, we have $[C] \in \mathbb{R}_{\geq 0} \beta$ by Lemma 2.9. By Lemma 2.6, we have $\dim W_2 \geq \mathcal{C}_{t_0} \cdot F_{t_0}^*(-K_{\mathcal{X}}) - 1 > \frac{n}{2}$. Since $W_1 \cap W_2$ is non-empty, it follows that

$$\dim(W_1 \cap W_2) \geq \dim W_1 + \dim W_2 - n \geq 1.$$

Therefore we can find a curve $C_1 \subset W_1 \cap W_2$. Again, we have $\mathbb{R}_{\geq 0} \alpha = \mathbb{R}_{\geq 0} \beta$.

Consider the extremal ray $\mathbb{R}_{\geq 0} \alpha$ and the corresponding extremal contraction $\phi_\alpha : X \rightarrow Y$. We will prove that Y is a point (and hence $\rho_X = 1$).

Since $\mathbb{R}_{\geq 0} \alpha = \mathbb{R}_{\geq 0} \beta$, for every $t \in T$ the curve $(\pi \circ F_t)(\mathcal{C}_t)$ is contracted by ϕ_α . Therefore $\bar{D} \subset \text{Exc}(\phi_\alpha)$. The morphism ϕ_α is either a divisorial contraction or a contraction of fiber type.

We first show that $\phi_\alpha : X \rightarrow Y$ can not be divisorial: Assume the contrary. Let A be a very ample divisor on Y and M a very ample divisor on X . Set

$$C' := D \cap_{i=1}^{\dim \phi_\alpha(D)} L_i \cap_{j=\dim \phi_\alpha(D)+1}^{n-2} M_j$$

where L_i 's are the general members of $|\phi_\alpha^* A|$ and M_j 's the general members of $|M|$. Let $S = \cap_{i=1}^{\dim \phi_\alpha(D)} L_i \cap_{j=\dim \phi_\alpha(D)+1}^{n-2} M_j$. Note that C' is contracted by ϕ_α . Therefore, it is a (positive) multiple of α . Note that $C' \cdot D = (C')_S^2 < 0$ since C' is a contracted curve on the surface S . Thus $\alpha \cdot D < 0$. This gives a contradiction.

Now we show that Y is a point. Assume by contradiction that $\dim Y > 0$. Let $y \in Y$ be a general point. The fiber X_y is a smooth Fano variety. Note that $K_X|_{X_y}$ is numerically equivalent to K_{X_y} . By Mori's bend and break, there is a curve $C' \subset X_y$ such that

$$C' \cdot (-K_{X_y}) \leq \dim X_y + 1.$$

Note that

$$C' \cdot (-K_{X_y}) = C' \cdot (-K_X) > \frac{n}{2} + 1$$

by our assumption. Combining these two inequalities yields $\dim X_y > \frac{n}{2}$. Thus $\dim Y < n - \frac{n}{2} = \frac{n}{2}$.

Let $\mathbb{R}_{\geq 0} \gamma$ be an extremal ray which is not contracted by ϕ_α . By Lemma 2.8, there is a twisted stable map $g : \mathcal{E}' \rightarrow \mathcal{X}$ such that (1) $\mathcal{E}'$ has at most 1 twisted point, (2) the intersection number $\mathcal{E}' \cdot (-K_{\mathcal{X}}) \leq n + 1$, and (3) $(\pi \circ g)_*[\mathcal{E}'] \in \mathbb{R}_{\geq 0} \gamma$. We may assume $(\pi \circ g)_*[\mathcal{E}'] = \gamma$.

Again, we view $g : \mathcal{E}' \rightarrow \mathcal{X}$ as an element of $\mathcal{K}_{0,1}(\mathcal{X}, \gamma)$ as in Section 2.4. Take an irreducible component $\mathcal{M} \subset \mathcal{K}_{0,1}(\mathcal{X}, \gamma)$ which contains $[g]$ and consider the universal family of twisted stable maps over $\mathcal{M}$ as in Section 2.4. Denote by W_3 the image of this family in X . For this family of twisted 1-pointed stable maps, the image of the marked point ∞ is also fixed.

Note that if $C'' \subset W_3$ is any curve, then its class is a multiple of γ by Lemma 2.9. By Lemma 2.6, we have $\dim W_3 > \frac{n}{2}$ and hence $\dim W_3 > \dim Y$. For any $y \in Y$, consider the intersection $W_3 \cap X_y$. It is easy to see that $\dim W_3 \cap X_y < 1$; otherwise, we would have $\mathbb{R}_{\geq 0} \alpha = \mathbb{R}_{\geq 0} \gamma$.

It follows that the morphism $\phi_\alpha|_{W_3} : W_3 \rightarrow Y$ is finite to its image. This is absurd since $\dim W_3 > \dim Y$. □

Notation 3.2. *Let*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{X} \\ G \downarrow & & \\ T & & \end{array}$$

be a covering family of twisted stable maps over an irreducible and projective scheme T . Pushing forward to the coarse spaces, we have

$$\begin{array}{ccc} C & \xrightarrow{\bar{F}} & X \\ \bar{G} \downarrow & & \\ T & & \end{array}$$

a covering family of stable maps into X . Let k be the number of twisted points on the source curve $\mathcal{C}_t$ for a general $t \in T$. Let $d = \mathcal{C}_t \cdot F_t^(-K_{\mathcal{X}})$. By Lemma 2.11, we may assume that $d \leq 2n$. Choose a covering family with the smallest possible anti-canonical degree. We fix this covering family of anti-canonical degree $d \leq 2n$ in Proposition 3.3 and Proposition 3.6.*

We proceed by dividing into two cases according to the number of twisted points on a general source curve: (1) $k \geq 2$, and (2) $k = 0$ or 1 . Proposition 3.3 takes care of the first case and Proposition 3.6 takes care of the second.

Case (1): $k \geq 2$.

Proposition 3.3. *Let $F : \mathcal{C} \rightarrow \mathcal{X}$ be the covering family of twisted stable maps as in Notation 3.2. Assume that $\mathcal{C}_t$ has at least two twisted points for a general $t \in T$. Also assume that $i_X > \max\{\frac{n}{2} + 1, \frac{2n}{3}\}$. Then $\rho_X = 1$.*

Remark 3.4. The condition $i_X > \frac{2n}{3}$ implies that the domain curve $\mathcal{C}_t$ has at most two components which are not contracted.

Proof. Let $S \subset T$ be any projective curve. Consider the pull-back family of twisted stable maps $F|_S : \mathcal{C}|_S \rightarrow \mathcal{X}$. Suppose that the image of this family

in $\mathcal{X}$ is two dimensional, i.e. the twisted curve deforms in $\mathcal{X}$. Recall that the images of twisted points are fixed in this family. By Mori's bend and break, this family has to degenerate.

Let $\tilde{D} \subset T$ be the loci where the domain curve is not irreducible. Consider $(\pi \circ F)(G^{-1}(\tilde{D})) \subset X$.

Claim 3.5. *The closed subset $(\pi \circ F)(G^{-1}(\tilde{D}))$ has at least one component of codimension at most 1.*

Proof of Claim 3.5. Assume by contradiction that every component of $(\pi \circ F)(G^{-1}(\tilde{D}))$ has codimension at least 2.

We can find a curve $E \subset X$ such that E does not meet $(\pi \circ F)(G^{-1}(\tilde{D}))$ and E is not contained in the image of any fiber of $\mathcal{C} \rightarrow T$ (by taking E the intersection of general ample divisors). Consider $G((\pi \circ F)^{-1}(E)) \subset T$. We can find a (complete) curve $E' \subset G^{-1}(G((\pi \circ F)^{-1}(E)))$ such that $F(E')$ is not contained in the image of any fiber of $\mathcal{C} \rightarrow T$. Take $E'' := G(E')$ and consider the family $F|_{E''} : \mathcal{C}_{E''} \rightarrow \mathcal{X}$. Note that the image of $F|_{E''} : \mathcal{C}_{E''} \rightarrow \mathcal{X}$ is two dimensional. By Mori's bend and break, this family has to degenerate, and hence E' has to meet $\tilde{D}$. This gives a contradiction. $\square$

Continue the proof of Proposition 3.3. Let D be an irreducible divisor contained in $(\pi \circ F)(G^{-1}(\tilde{D}))$. Consider possible degenerations of twisted stable maps. By degree reason, only two components of $\mathcal{C}_t$ are not contracted by F_t . Denote these two components by $\mathcal{C}_t^1$ and $\mathcal{C}_t^2$. Note that $(\pi \circ F_t)(\mathcal{C}_t^1)$ and $(\pi \circ F_t)(\mathcal{C}_t^2)$ are unsplit. If $\mathcal{C}_t^i$, $i = 1, 2$, has two or more twisted points, then its image in $\mathcal{X}$ can not deform by Mori's bend and break argument.

Since the locus of degenerate curves in X contains at least a divisor, it follows that at least one component of the domain curve, say $\mathcal{C}_t^1$, will have at most one twisted point. Thus there is a family of twisted stable maps with at most one twisted point and its image in X contains at least a divisor. We apply Lemma 3.1 and conclude the proof of Proposition 3.3. $\square$

Case (2): $k \leq 1$.

Proposition 3.6. *Let $F : \mathcal{C} \rightarrow \mathcal{X}$ be the covering family as in Notation 3.2. Assume that $\mathcal{C}_t$ has at most one twisted point for a general $t \in T$. Also assume that $i_X > \max\{\frac{n}{2} + 1, \frac{2n}{3}\}$. Then $\rho_X = 1$.*

Proof. We first prove the following claim:

Claim 3.7. *For a very general point $x \in \mathcal{X}$, there is a twisted rational curve $\mathcal{E}$ through x such that (1) $\mathcal{E}$ has at most one twisted point, and (2) $\mathcal{E}$ has $-K_{\mathcal{X}}$ -degree at most $n + 1$.*

Proof of Claim 3.7. Note that there are only countably many unsplit classes $\alpha_i \in H_2(X)$, $i \in I$. Consider all possible twisted stable maps such that (1) the source curve is birational to its image, (2) the source curve has at least two twisted points, and (3) the class of the image is one of these α_i 's. For each α_i there are only finitely many such twisted stable maps thanks to

Mori's bend and break. Delete the images of all such twisted stable maps and all twisted points from $\mathcal{X}$. Denote the resulting set by U .

Let $x \in U$. Pick a $t \in T$ such that x lies on the image of $\mathcal{C}_t$ (under the morphism F_t). Note that x is the image of an ordinary point, denoted by $0 \in \mathcal{C}_t$, since the image of a twisted point is a twisted point. If $\mathcal{C}_t$ has a twisted point, we denote it by ∞ . If $\mathcal{C}$ does not have any twisted point, pick any point on $\mathcal{C}_t - \{0\}$ whose image in $\mathcal{X}$ is not x . Also denote this point by $\infty \in \mathcal{C}_t$.

Suppose that $\mathcal{C}_t \cdot F_t^*(-K_{\mathcal{X}}) > n + 1$. Then

$$\dim_{[F_t]} \text{Mor}(\mathcal{C}_t, \mathcal{X}, F_t|_{\{0, \infty\}}) > 1$$

by Lemma 2.2.

By Mori's bend and break, the family $F : \mathcal{C} \rightarrow \mathcal{X}$ degenerates at some $t' \in T$. By degree reason, there are only two components of $\mathcal{C}_{t'}$, say $\mathcal{C}_{t'}^1$ and $\mathcal{C}_{t'}^2$, which are not contracted by $F_{t'}$. Note that since $F_{t'}$ is a twisted stable map, every contracted component needs to have at least three special points (nodes or the original twisted point). It implies that the domain curve $\mathcal{C}_{t'}$ has either two or three components.

In the first case, the two components $\mathcal{C}_{t'}^1$ and $\mathcal{C}_{t'}^2$ intersect at a node which may be a new twisted point. In the second case, there are three components $\mathcal{C}_{t'}^1$, $\mathcal{C}_{t'}^2$ and $\mathcal{C}_{t'}^3$. The contracted component $\mathcal{C}_{t'}^3$ has three special points: the original twisted point ∞ and two nodes $p_1 = \mathcal{C}_{t'}^1 \cap \mathcal{C}_{t'}^3$ and $p_2 = \mathcal{C}_{t'}^2 \cap \mathcal{C}_{t'}^3$. Note that p_1 and p_2 may be new twisted points. By this analysis, we can obtain a curve, say $F_{t'} : \mathcal{C}_{t'}^1 \rightarrow \mathcal{X}$, such that the image $F_{t'}(\mathcal{C}_{t'}^1)$ contains x and the source curve $\mathcal{C}_{t'}^1$ has at most two twisted points (the original twisted point ∞ and the possibly new twisted point at the node). We claim that $\mathcal{C}_{t'}^1$ can not have two twisted points: Suppose that $\mathcal{C}_{t'}^1$ does have two twisted points. The image of $\mathcal{C}_{t'}^1$ (under $\pi \circ F_{t'}$) will be in $X - U$. This is not possible since $x \in U$.

Therefore we obtain a twisted rational curve through x with at most 1 twisted point and of smaller anti-canonical degree. Repeat this bend and break process until the anti-canonical degree is at most $n + 1$. This concludes the proof of Claim 3.7. $\square$

Now we show that the covering family $F : \mathcal{C} \rightarrow \mathcal{X}$ has anti-canonical degree $d \leq n + 1$.

Assume the contrary, i.e. $d > n + 1$. Let $r \in \mathbb{N}$ be the natural number such that rK_X is a Cartier divisor as in Notation 2.7. Note that $r(C \cdot -K_X) \in \mathbb{N}_{>0}$ for any curve $C \subset X$. Let l be any rational number such that $0 < l \leq n + 1$ and $rl \in \mathbb{N}_{>0}$. Consider $\mathcal{K}_{0,1}(\mathcal{X}, l)$ and its universal family of twisted stable maps $F_l : \mathcal{U}_l \rightarrow \mathcal{X}$. The union $\cup_{\{l|0 < l \leq n+1, rl \in \mathbb{N}\}} (\pi \circ F_l)(\mathcal{U}_l)$ is a closed set since it is the union of finitely many closed set. It is clear that $\cup_{\{l|0 < l \leq n+1, rl \in \mathbb{N}\}} (\pi \circ F_l)(\mathcal{U}_l) = X$ since through any very general point $x \in \mathcal{X}$ there is a twisted rational curve with at most 1 twisted point and of anti-canonical degree $l \leq n + 1 < d$ by Claim 3.7. Since X is Noetherian and

irreducible, we have $(\pi \circ F_l)(\mathcal{U}_l) = X$ for some l . By the same argument in Lemma 2.11, we have a covering family of degree strictly less than d over an irreducible and projective scheme. This is not possible since our choice of the covering family $F : \mathcal{C} \rightarrow \mathcal{X}$ is of the smallest (anti-canonical) degree. Thus we have $d \leq n + 1$.

Note that $d \leq n + 1$ implies that the covering family is unsplit by degree reason. Now apply Lemma 3.1 to conclude the proof. $\square$

Proof of Theorem 1.4. Theorem 1.4 now follows easily from Proposition 3.3 and Proposition 3.6. $\square$

The statement

$$i_X > \frac{n}{2} + 1 \Rightarrow \rho_X = 1$$

is equivalent to

$$\rho_X \geq 2 \Rightarrow i_X \leq \frac{n}{2} + 1.$$

Under an extra assumption, we are able to prove the following:

Proposition 3.8. *Let X be a Fano variety with at worst isolated quotient singularities. Assume that $\rho_X \geq 2$ and there is an extremal contraction $\phi : X \rightarrow Y$ such that $\dim \text{Exc}(\phi) \geq n - 1$. Then $i_X \leq \frac{n}{2} + 1$.*

Proof. Assume the contrary that $i_X > \frac{n}{2} + 1$. Divide into two cases: (1) the contraction $\phi : X \rightarrow Y$ is divisorial, and (2) the contraction is of fiber type.

First consider the divisorial case. Let $D = \text{Exc}(\phi) \subset X$ be the exceptional divisor, and $\mathbb{R}_{\geq 0} \alpha$ be the contracted extremal ray. As in the proof of Lemma 3.1, we can find an extremal ray $\mathbb{R}_{\geq 0} \beta$ such that $\beta \cdot D > 0$. Let $g : \mathcal{C} \rightarrow \mathcal{X}$ be a twisted stable map such that $\mathcal{C}$ has at most one twisted point and the twisted stable map $g : \mathcal{C} \rightarrow \mathcal{X}$ is birational to its image. We may also assume $[(\pi \circ g)(\mathcal{C})] = \beta$. Consider $g : \mathcal{C} \rightarrow \mathcal{X}$ as a twisted 1-pointed stable map of genus 0 with homology class β . Let $[g] \in \mathcal{M} \subset \mathcal{K}_{0,1}(\mathcal{X}, \beta, g|_{\infty})$ be an irreducible component. Consider its universal family of twisted 1-pointed stable maps $G : \mathcal{U} \rightarrow \mathcal{X}$. Denote by $W \subset X$ the image of this family in X . Note that $\dim W > \frac{n}{2}$ (by Lemma 2.6). Take an irreducible component of $W' \subset W \cap D$. We have $\dim W' > \frac{n}{2} + (n - 1) - n = \frac{n}{2} - 1$. Note that for any curve $C' \subset W' \subset W$, its class $[C']$ lies in the ray $\mathbb{R}_{\geq 0} \beta$ (by Lemma 2.9).

Set $Z = \phi(D)$ (with the reduced scheme structure). Consider $\phi|_D : D \rightarrow \phi(D) = Z$. By a similar argument (using intersection number) as in the proof of Lemma 3.1, we have $\mathbb{R}_{\geq 0} \beta \neq \mathbb{R}_{\geq 0} \alpha$.

It is easy to see that $\dim Z > 0$: If $\dim Z = 0$, then $\mathbb{R}_{\geq 0} \alpha = \mathbb{R}_{\geq 0} \beta$; a contradiction. Pick a general $y \in Z \subset Y$ and consider the fiber X_y . Since X has only isolated singularities, we may assume that X_y is smooth (and the morphism ϕ is smooth near X_y). Since the exceptional divisor D is covered by rational curves in the ray $\mathbb{R}_{\geq 0} \alpha$, there is a (twisted) stable map $h : \mathbb{P}^1 \rightarrow X_y \subset \mathcal{X}$ such that $[(\pi \circ h)(\mathbb{P}^1)] \in \mathbb{R}_{\geq 0} \alpha$. We may assume that $h : \mathbb{P}^1 \rightarrow X_y \subset \mathcal{X}$ is birational to its image. If $\mathbb{P}^1 \cdot h^*(-K_X) > n + 1$, it is easy

to see that $\dim \operatorname{Mor}_{[h]}(\mathbb{P}^1, \mathcal{X}, h|_{\{0, \infty\}}) > 1$. Applying bend and break, we obtain a rational curve with smaller degree. Note that the resulting rational curve is still on X_y and its class is a multiple of α (since we start with a curve whose class is in the extremal ray $\mathbb{R}_{\geq 0} \alpha$). Continuing this bend and break process, we may assume that $h : \mathbb{P}^1 \rightarrow X_y \subset \mathcal{X}$ has $-K_{\mathcal{X}}$ -degree at most $n + 1$. Hence it is unsplit.

As before, we can find an irreducible component $[h] \in \mathcal{M}_1 \subset \mathcal{K}_{0,1}(\mathcal{X}, \alpha)$ and its universal family of twisted 1-pointed stable maps $H : \mathcal{U}_1 \rightarrow \mathcal{X}$. Denote by W_1 its image in X . Note that $W_1 \subset X_y$. Therefore, we have $\dim X_y \geq \dim W_1 > \frac{n}{2}$. Also note that $\phi|_{W'} : W' \rightarrow Z$ is finite to its image. Therefore, we have $\dim Z \geq \dim W'$. It follows that $\dim D = \dim X_y + \dim Z > \frac{n}{2} + \frac{n}{2} - 1 = n - 1$. This gives a contradiction.

Now consider the fiber type contraction case. The proof is quite similar to the proof of the fiber type case in Lemma 3.1. Let $\mathbb{R}_{\geq 0} \beta$ be an extremal ray not contracted by the morphism ϕ . We can find a twisted 1-pointed stable map $h : \mathcal{C} \rightarrow \mathcal{X}$ of genus 0 and homology class β . Let $[h] \in \mathcal{M}_2 \subset \mathcal{K}_{0,1}(\mathcal{X}, \beta, h|_{\infty})$ be an irreducible component and $H : \mathcal{U}_2 \rightarrow \mathcal{X}$ its universal family of twisted 1-pointed stable maps. Denote by W_2 the image of $\mathcal{U}_2$ in X . Let $y \in Y$ be a general point and X_y be the fiber. As in the proof of Lemma 3.1, we have $\dim X_y > \frac{n}{2}$ and $\dim Y = \dim X - \dim X_y < \frac{n}{2}$. Note that $\dim W_2 > \frac{n}{2}$ (Lemma 2.6) and the morphism $\phi|_{W_2} : W_2 \rightarrow Y$ is finite to its image. Hence $\dim Y \geq \dim W_2 > \frac{n}{2}$. This gives a contradiction. $\square$

REFERENCES

- [ACV03] D. Abramovich, A. Corti, and A. Vistoli, *Twisted bundles and admissible covers*, Comm. Algebra 31 (2003) 3547–3618.
- [AV02] D. Abramovich, A. Vistoli. *Compactifying the space of stable maps*, J. Amer. Math. Soc. 15, no. 1 27–75, 2002.
- [ACO04] M. Andreatta, E. Chierici, and G. Occhetta. *Generalized Mukai Conjecture for Special Fano Varieties*, Cent. Eur. J. Math. 2 (2004), no. 2, 272–293, 2004.
- [BCDD03] L. Bonavero, C. Casagrande, O. Debarre and S. Druel. *Sur une conjecture de Mukai*, Comment. Math. Helv. 78 (2003), no. 3, 601–626, 2003.
- [CMSB02] K. Cho, Y. Miyaoka and N.I. Shepherd-Barron. *Characterizations of projective spaces and applications to complex symplectic manifolds*, Higher dimensional birational geometry (Kyoto, 1997), 1–88, Adv. Stud. Pure Math., 35, Math. Soc. Japan, Tokyo, 2002.
- [Ch06] J.-C. Chen. *Characterizing projective spaces for varieties with at most quotient singularities*, Preprint math.AG/0604522, 2006.
- [CT09] J.-C. Chen and H.-H. Tseng. *Cone Theorem via Deligne-Mumford stacks*, Preprint math.AG/0505043, to appear in Math. Ann. 2009.
- [CT07] J.-C. Chen and H.-H. Tseng. *Note on Characterization of Projective spaces*, Comm. Algebra 35 (2007), no. 11, 3808–3819, 2007.
- [De01] O. Debarre. *Higher-dimensional algebraic geometry*, Springer-Verlag, 2001.
- [Ka91] Y. Kawamata, *On the length of an extremal rational curve*, Invent. Math. 105 (1991), no. 3, 609–611.

- [Ke01] S. Kebekus. *Characterizing the projective space after Cho, Miyaoka and Shepherd-Barron*, Complex Geometry (Göttingen, 2000)147-155, Springer, Berlin, 2002.
- [Ko96] J. Kollár. *Rational curves on algebraic varieties*, Springer-Verlag, 1996.
- [KM98] J. Kollár, S. Mori. *Birational Geometry of Algebraic Varieties*, Cambridge Tracts in Mathematics 134, Cambridge University Press, 1998.
- [Mu88] S. Mukai. *Open problems*, In The Taniguchi foundation, editor, Birational geometry of algebraic varieties., Katata, 1988.
- [Wi90] J. A. Wiśniewski. *On a conjecture of Mukai*, Manuscripta Math. 68 (1990), 135141.

NATIONAL CENTER FOR THEORETICAL SCIENCES,, MATH. DIVISION, THIRD GENERAL BUILDING, NATIONAL TSING HUA UNIVERSITY, NO. 101 SEC 2 KUANG FU ROAD, HSINCHU, TAIWAN 30043, TAIWAN

E-mail address: `jcchen@math.nthu.edu`, `jcchenster@gmail.com`